



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

115. Proposed by MARY BLAINE, B. Sc., Graduate Student, Drury College, Springfield, Mo.

The locus of a point such that the sum of the squares of its normals form a given ellipsoid is constant, is a co-axial ellipsoid. [From *C. Smith's Solid Analytical Geometry*, page 95.]

*** Solutions of these problems should be sent to B. F. Finkel, not later than March 10.

CALCULUS.

85. Proposed by J. SCHEFFER, A. M., Hagerstown, Md.

A line of double curvature, beginning at some point in the circumference of the base circle of a right cone, winds itself under the constant inclination β to the base circle around the curved surface of the cone. Find its length and that of its projection upon the base circle.

86. Proposed by WALTER H. DRANE, Graduate Student, Harvard University, 65 Hammond Street, Cambridge, Mass.

Prove that the curve whose normal equals its radius of curvature drawn in an opposite direction, is the catenary, $y = c \cosh(x/c)$.

87. Proposed by MARY BLAINE, B. Sc., Graduate Student, Drury College, Springfield, Mo.

Integrate $(px - y)(py + x) = h^2 p$, where $p = dy/dx$.

*** Solutions of these problems should be sent to J. M. Colaw, not later than March 10.

MECHANICS.

82. Proposed by WALTER H. DRANE, Graduate Student, Harvard University, 65 Hammond Street, Cambridge, Mass.

A sphere, diameter $2a$, rests in limiting equilibrium upon the edge of a box and against a vertical wall. If the box be of such dimensions that it will not tip, find the distance of the box from the wall, having given the coefficient of friction between the sphere and wall $\frac{1}{2}$, between the sphere and box $\frac{1}{3}$, and between the box and floor $\frac{1}{3}$. [From *Problems in Mechanics* proposed to class in Harvard University.]

83. Proposed by MARY BLAINE, B. Sc., Graduate Student, Drury College, Springfield, Mo.

A particle is projected upwards in vacuo with a velocity v . Show that on reaching the ground again there is no deviation to the south, but the deviation to the west is $4\omega \cos \lambda (v^3/3g^2)$. [*Laplace*, iv, page 341.]

*** Solutions of these problems should be sent to B. F. Finkel, not later than March 10.

DIOPHANTINE ANALYSIS.

78. Proposed by COOPER D. SCHMITT, A. M., Professor of Mathematics, University of Tennessee, Knoxville, Tenn.

Find three square numbers in harmonical progression.

79. Proposed by EDMUND FISH, Hillsboro, Ill.

Find an integral right triangle in which the bisector of one of the acute angles is also integral.

*** Solutions of these problems should be sent to J. M. Colaw, not later than March 10.